arXiv:1109.1942v1 [nucl-th] 9 Sep 2011

# Doppler effect and angular correlation for the precision measurement of $^{29}$Si binding energy

Yongkyu Ko[a], and K. S. Kim[b]

*Research Institute for Basic Sciences,*

*Korea Aerospace University, Koyang 412-791, Korea*

(Dated: September 12, 2011)

The importance of the Doppler effect of the recoiling intermediate nucleus in cascade decay, for an example, $^{29}$Si, is emphasized in the precision measurement of the nuclear binding energy using a flat-crystal spectrometer. A more accurate experiment method that a large uncertainty due to the Doppler effect of the recoiling nucleus could be eliminated by a coincidence measurement of two successive $\gamma$-rays taken into account their angular correlation is suggested in the framework of relativistic kinematics. It is also suggested that this method can apply to the nuclear lifetime measurement of intermediate states.

PACS numbers: 21.10.Dr, 25.40.Lw, 27.30.+t, 29.30.Kv

---

[a] yongkyu.ko@gmail.com

[b] kyungsik@kau.ac.kr

Nuclear binding energy is of great importance as shown in the first chapter of nuclear physics and it is indispensable for studying the structure of a nucleus. Recently binding energies are accurately measured by two kinds of method. one of them is a $\gamma$-ray measurement in a neutron capture reaction by using a flat-crystal spectrometer [1], and the other is the mass comparison of two different ions simultaneously confined in a Penning trap by measuring their cyclotron frequencies [2, 3]. The comparison of the two experiments for $^{29}$Si and $^{33}$S was reported in association with Einstein's famous formula $E = mc^2$, and the systematic error associated with either measurement was given; the error on the comparison is currently dominated by the uncertainty on the $\gamma$-ray measurements [4]. It is also reported in Ref. [1] that the uncertainty for $^{29}$Si can be shown from the significantly Doppler broadened 4934-keV profile, which decreases the accuracy with which the Bragg angle can be determined. The 4934-keV line is broadened by the Doppler effect, because it is emitted while the nucleus is recoiling following the emission of the 3539-keV line.

In an accurate $\gamma$-ray measurement like the flat-crystal spectrometer, the importance of the Doppler effect caused by the incident neutron was pointed out in the determination of deuteron binding energy [5]. So the effect was considered in Ref. [1], but was reported to be negligible. It is evident that the Doppler effect in case of heavy nuclei does not so significant as in case of the deuteron [5], because it depend on the reciprocal of the target mass. In cascade decay, however, the recoil velocity of the intermediate nucleus caused by the emission of the primary $\gamma$-ray is significantly large(39 km/s) for the $^{29}$Si, and thus the Doppler shift of the secondary $\gamma$-ray energy is considerable($-646.9 \cos\theta$ eV) according to the direction of a measurement. If we were able to detect the secondary $\gamma$-ray at a specified angle, a precision measurement could be achieved. But the secondary $\gamma$-rays are measured from the sources which are in random motion, so that the spectrum of the $\gamma$-rays is broadened, because all the Doppler shifted $\gamma$-rays emitted from the sources are summed in the spectrum. They do not vanish as explained in Ref. [1]. To solve this problem in Ref. [1], the number of measurements for the 4934-keV line is increased to 147 times, while the number of measurements for the 3539-keV line is only 42, since the statistic error is weighted by the inverse of the square root of the number of measurement.

On the other hand, a coincidence measurement using angular correlation in cascade decay would be a good solution to the precision measurement [6]. The secondary $\gamma$-ray has angular correlation with respect to the primary $\gamma$-ray, if the two $\gamma$-rays are simultaneously measured.

The most probable directions of the secondary $\gamma$-ray are parallel and anti-parallel with the direction of the primary $\gamma$-ray, and the Doppler shift of the secondary $\gamma$-ray is maximum at these angles. Therefore, the average value of the two energies of the Doppler shifted $\gamma$-rays at angle 0 and 180 is free from Doppler broadening caused by the recoil of the nucleus, and thus the profile of the shifted $\gamma$-ray might be even closer to a natural line width which can also give the lifetime of the intermediate state of the nucleus. The Institut Laue-Lagnevin (ILL) high-flux reactor has a through beam tube [7], and it is possible to perform a coincidence measurement. A difficulty may arise in this experiment, that is, the experimental run time should be increased. It can be a solution that increasing the amount of the target source can increase its activity within the allowed solid angle of the detector. Unfortunately, this problem is beyond the scope of our discussion, and our propose is to give a suggestion to improve a precision measurement from the Doppler effect and angular correlation in the framework of relativistic kinematics. A precision experiment taken into account the Doppler effect and angular correlation provides a measurement of the lifetime of intermediate states as well as a measurement of accurate binding energies, because the angle dependent measurements of the secondary $\gamma$-ray give the information of the attenuated velocity of the recoiling nucleus, and one can expect to see the secondary $\gamma$-ray profile which is comparable to the natural line width of an intermediate state.

The binding energy for a neutron capture reaction is the mass difference between initial and final states as follows:

$$m(n) + m(^{A}X) = m(^{A+1}X) + S_n, \tag{1}$$

where $m$ means the mass of the particle in parentheses, $A$ is an atomic number, and $S_n$ is the separation energy of a neutron or the binding energy of a neutron which is carried by $\gamma$-rays and the nucleus. If it is a dominant channel that a single $\gamma$-ray is emitted in a neutron capture reaction, the problem is simple and has been already considered in Ref. [5]. In this letter we are interested in the decay with emitting two $\gamma$-rays successively which is the most probable channel. The excited state of $^{29}$Si is the case as shown in Fig. 1 and the values in parentheses are the number of $\gamma$-rays emitted per 100 neutron captures [8]. This decay channel can be represented as a Feynman diagram shown in Fig. 2 and can be separated into two parts as Figs. 3, and 4. The equation of the binding energy Eq. (1) is also separated into two parts corresponding to the separated Feynman diagrams, respectively, as follows:

$$m(n) + m({}^{A}X) = m({}^{A+1}X^{*}) + E_{b1}, \tag{2}$$

$$m({}^{A+1}X^{*}) = m({}^{A+1}X) + E_{b2}, \tag{3}$$

$$S_n = E_{b1} + E_{b2}, \tag{4}$$

where asterisk means an intermediate state of the final nucleus, namely, an excited state of the final nucleus.

Fig. 3 represents the neutron capture reaction ${}^{28}\mathrm{Si}(n,\gamma){}^{29}\mathrm{Si}^{*}$ and the energy-momentum conservation law is given by

$$m + M = E_1 + \omega_1, \tag{5}$$

$$0 = \boldsymbol{p}_1 + \boldsymbol{k}_1, \tag{6}$$

where $m = m(n)$ and $M = m({}^{28}\mathrm{Si})$. The energy-momentum relations of the final state are $E_1^2 = \boldsymbol{p}_1^2 + M_1^2$ for the intermediate state of the silicon nucleus, and $\omega_1^2 = \boldsymbol{k}_1^2$ for the primary photon. It should be noted that the velocity of the center of mass for the neutron and the silicon system is only 114 $m/s$ for the incident neutron flux energy 0.056 eV, while that for the neutron and the proton system is 1.6 $km/s$ as calculated in Ref. [5]. Hence the Doppler effect of the silicon nucleus due to the incident neutron is negligible ($-1.3\cos\theta$ eV) so that it is not necessary to consider the kinetic energy of the incident neutron.

Solving the equations of the energy-momentum conservation laws with respect to the primary photon energy and replacing the initial masses by the binding energy, one can obtain the binding energy of the intermediate nucleus as follows:

$$E_{b1} = \omega_1 - M_1 + \sqrt{\omega_1^2 + M_1^2} \cong \omega_1 + \frac{\omega_1^2}{2M_1}, \tag{7}$$

which is reduced to the non-relativistic result that are usually used in the literature. Since the incident kinetic energy of the neutron is ignored, there is no Doppler effect in this equation. The last term in the above equation is the kinetic energy of the recoiling intermediate nucleus, and the velocity of it can be checked as follows:

$$KE = \frac{1}{2} M_1 v_{int}^2 = \frac{\omega_1^2}{2M_1}, \;\; v_{int} \cong 39 km/s, \tag{8}$$

where the natural unit($\hbar = c = 1$) is used in every equation throughout this letter, and thus the velocity in SI unit should be multiplied by the light velocity.

Since this velocity causes the secondary $\gamma$-ray to be shifted considerably, it is important to take into account the kinetic energy and momentum of the intermediate nucleus in calculation. Hence the law of the energy-momentum conservation for the process shown in Fig. 4 should be given by

$$E_1 = E_2 + \omega_2, \tag{9}$$

$$\boldsymbol{p}_1^2 = \boldsymbol{p}_2^2 + \boldsymbol{k}_2^2, \tag{10}$$

where the energy-momentum relation of the ground state silicon nucleus is $E_2^2 = \boldsymbol{p}_2^2 + M_2^2$, and that of the secondary photon is $\omega_2^2 = \boldsymbol{k}_2^2$. To specify the angle between two photons that is of importance in the Doppler effect and angular correlation, it is useful to show the following intermediate step:

$$(E_1 - \omega_2)^2 = E_2^2 = M_2^2 + (\boldsymbol{p}_1^2 + \boldsymbol{k}_2^2 - 2p_1k_2\cos\theta), \tag{11}$$

where the direction of the primary photon is $\theta = \pi$ and the direction of the recoiling nucleus is $\theta = 0$. Solving the equation as the same way of the previous energy-momentum law, one can obtain the binding energy of the ground state of the silicon as follows:

$$\begin{aligned} E_{b2} &= \omega_2 - M_2 + \sqrt{M_2^2 + \omega_2^2 + 2\omega_2\{E_{b1} - \omega_1(1+\cos\theta)\}} \\ &\cong \omega_2 + \frac{\omega_2^2}{2M_2} + \frac{\omega_1^2\omega_2}{2M_1M_2} - \frac{\omega_1\omega_2}{M_2}\cos\theta, \end{aligned} \tag{12}$$

where the second term is the kinetic energy due to recoil, the third term is negligible kinetic term, and the last term represents the Doppler effect as explained in Ref. [5]. This result is calculated in laboratory frame and is consistent with the result that is Lorentz transformed from the calculations in rest frame of $^{29}$Si$^*$.

The total binding energy of the silicon is the sum of Eqs. (7) and (12) as follows:

$$\begin{aligned} S_n = E_{b1} + E_{b2} &\cong \omega_1 + \omega_2 + \frac{\omega_1^2}{2M_1} + \frac{\omega_2^2}{2M_2} + \frac{\omega_1^2\omega_2}{2M_1M_2} - \frac{\omega_1\omega_2}{M_2}\cos\theta \\ &= 3538966.3(1.6) + 4933946.3(4.0) + 232.9 + 450.9 + 0.0 - 646.9\cos\theta \text{ eV}. \end{aligned} \tag{13}$$

If one is able to handle the angle in the secondary $\gamma$-ray measurement, the Doppler shifted $\gamma$-ray is detected. Otherwise, all the shifted $\gamma$-rays reach the detector and appear in the spectrum as a broadened lineshape. The Doppler effect of the secondary $\gamma$-ray is so large that the precision measurement of the flat-crystal spectrometer can be questionable without considering the effect. Increasing the number of measurements may not reduce the error fundamentally. The angular correlation method which has long been established [6] can be applied to this cascade decay measurement to reduce the uncertainty broadened by the Doppler effect of the recoiling nucleus. Fig. 5 shows the idea of a coincidence measurement. If the two $\gamma$-rays are measured simultaneously, the secondary $\gamma$-ray has angular correlation in association with the primary $\gamma$-ray, namely, the distribution of the secondary $\gamma$-ray with respect to the primary $\gamma$-ray varies from angle to angle.

Since the polarizations of the $\gamma$-rays are not observed and only directional correlation is observed, the angular correlation function [6, 10] is simply given by

$$W(\vartheta) = \sum_{l=even}^{l_{max}} A_l P_l(\cos\vartheta), \tag{14}$$

where $P_l$ is the Legendre polynomial of order $l$. The coefficient for the transition of angular momentum states $J_A \to J_B \to J_C$ with emitting photons of the angular momenta $L_1$ and $L_2$ in order is given by

$$A_l = F_l(L_1, J_A, J_B) F_l(L_2, J_C, J_B), \tag{15}$$

$$F_l = W(J_B J_A l L; L J_B) C_{l0}(LL), \tag{16}$$

where $W(J_B J_A l L; L J_B)$ is a Racah coefficient, and $C_{l0}(LL) \propto (-1)^{L-1}(2L+1) < L, L, 1, -1|l0 >$ is known as the radiation parameter. The angle of the binding energy in Eq. (13) is related with the angle in the correlation function as $\theta = \pi - \vartheta$, but it is not necessary to distinguish between the angles, because only even power of the Legendre polynomial contributes to the angular correlation function. For a simple example, if a dipole-dipole transition occurs through the nuclear states of $J_A = J_C = 0$, and $J_B = 1$, the angular correlation function is $W(\vartheta) = \frac{3}{16\pi}(1 + \cos^2\vartheta)$. If a dipole-dipole transition takes place through the nuclear states of $J_A = J_C = 1/2$ , and $J_B = 3/2$ as assigned from the $\gamma$-rays measurement in Ref. [9] and shown in Fig. 1, the angular correlation function is

$$W(\vartheta) = \frac{3}{32\pi}(1 + \frac{3}{7}\cos^2\vartheta). \tag{17}$$

One can see here that the secondary $\gamma$-ray has the most probable distribution at the angles $\theta = 0$, or $\pi$, and it is shifted maximally according to Eq. (13) because of the Doppler effect. The best coincidence measurement of the two $\gamma$-rays would be at the angle $\vartheta = \pi$, that is, the two $\gamma$-rays are radiating opposite each other.

However, there is a problem that the surrounding environment in which the recoiling nucleus is located is not a free space. Therefore, it is not sure that the shifted $\gamma$-ray at the angle $\theta = 0$ gives exact information on the binding energy. Moreover the surrounding environment varies from nucleus to nucleus. For an example, the velocities of the intermediate state of other nuclei $^{33}$S and $^{36}$Cl in Ref. [1] are greater than that of $^{29}$Si, because their primary $\gamma$-ray energies are larger than that of $^{29}$Si for their masses. However, there is no mention that the secondary $\gamma$-rays in the nuclei $^{33}$S and $^{36}$Cl suffer from the Doppler broadening as in case of $^{29}$Si. In general, the angular correlation of two $\gamma$-rays emitting successively is evident, when the lifetime of the intermediate state is sufficiently short [10]. The case also means that the Doppler effect is evident, because the recoil velocity of the intermediate nucleus is not much slowed down by interacting with the surrounding environment. Hence the lifetime of the intermediate state of $^{33}$S and $^{36}$Cl is much longer than that of $^{29}$Si as shown from experimental data [11]. The velocity of these intermediate nuclei could be attenuated during their long lifetimes, and thus the Doppler broadening would be considerably reduced. This aspect also reflects in the branching ratio of the decay, that is, the number of $\gamma$-ray per 100 neutron capture in Fig. 1 of Ref. [1]. Since the lifetime of the $^{29}$Si intermediate nucleus is quite short, and it is likely that the velocity of the intermediate nucleus is a little slowed down. However it is not sufficiently satisfactory to extract the exact binding energy from the above single coincidence measurement at an angle.

By using the Doppler broadening of the secondary $\gamma$-ray and the slowing down mechanism of the recoil nucleus, the lifetime of the intermediate state has been measured by the methods of the Doppler shift attenuation [12] and the gamma ray induced Doppler broadening [13]. In order to take into account the slowing down of the nucleus, these methods use several models, such as, the mean-free-path approach, molecular dynamics, and the phonon creation model. However, if a coincidence measurement with angular correlation is performed twice at angle $\theta = 0$ and angle $\theta = \pi$, one can measure the binding energy and the lifetime of the intermediate state quite simply. Though the velocity of the recoiling nucleus is attenuated by the slowing down process, the magnitude of the Doppler shifted secondary $\gamma$-ray energies can be exactly determined by the comparison of two coincidence measurements at angle $\theta = 0$ and $\pi$. The average value of the two shifted $\gamma$-ray energies is free from the Doppler broadening and the slowing down mechanism of the recoiling nucleus as follows:

$$\begin{aligned} S_n &= \frac{1}{2}\{S_n(\theta = 0) + S_n(\theta = \pi)\}, \\ \Delta E &= \frac{1}{2}\{S_n(\theta = 0) - S_n(\theta = \pi)\}, \end{aligned} \tag{18}$$

where the first equation is the desired binding energy, and the second equation is the magnitude of the Doppler shift attenuated by the surrounding environment. If this magnitude is compared to the calculated value in Eq. (13), one can extract the attenuated velocity of the recoiling nucleus which provides a lifetime in the Doppler shift attenuation method. A flat-crystal spectrometer could resolve the attenuated velocity, even though a short lifetime gives a little attenuation. Therefore, the uncertainty due to the nucleus recoil can be ruled out in the determination of the binding energy. In addition, since the line shape of the Doppler shifted $\gamma$-ray does not include the major broadening by the recoil of the nucleus but contains the minor broadening by the thermal motion in the lattice of the target, the decay width of the intermediate nucleus is much more comparable to the thermal broadened line shape. Hence, it is possible to compare the lifetime extracted from the magnitude of the Doppler shift with the thermal broadened linewidth which can be obtained from the classical Maxwell-Boltzman distribution by virtue of the modified Lorenzian shape used in Ref. [13].

In conclusion, the Doppler shift due to the velocity of the intermediate nucleus of $^{29}$Si is so large that a precision measurement of the secondary $\gamma$-ray should take into account the Doppler effect. A coincidence measurement of two $\gamma$-rays with considering their angular correlation can reduce the uncertainty caused by the Doppler effect of the recoiling nucleus in the determination of binding energy. This improved binding energy measurement would lead to one step upgrade for the direct test of $E = mc^2$ [4]. If the measurement technique for cascade decay is developed, one can learn the more accurate information about the nuclear structure, for examples, accurate energy level, the lifetime of the nuclear intermediate state, and so on.

## ACKNOWLEDGEMENTS

This work was supported by the Korea Science and Engineering Foundation(KOSEF) grant funded by the Korea government(MOST) (No. R01-2007-000-20569-0).

---

[1] M. S. Dewey, E. G. Kessler Jr., R. D. Deslattes, H. G. Börner, M. Jentshel, C. Doll, and P. Mutti, Phys. Rev. **C 73**, 044303 (2006).

[2] W. Shi, M. Redshaw, and E. G. Myers, Phys. Rev. **A 72**, 022510 (2005).

[3] S. Rainville, J. K. Thmpson and D. E. Pritchard, Science **303**, 334 (2004).

[4] S. Rainville, *et al.*, Nature(London) **438**, 1096 (2005).

[5] Y. Ko, M. K. Cheoun, and I. T. Cheon, Phys. Rev. **C 59**, 3473 (1999).

[6] L. C. Biedenharn and M. E. Rose, Rev. Mod. Phys. **25**, 729 (1953).

[7] E. G. Kessler Jr., M. S. Dewey, R. D. Deslattes, A. Henins, H. G. Börner, M. Jenschel, C. Doll, and H. Lehmann, Nucl. Instrum. Methods Phys. Res. **A 457**, 187 (2001).

[8] M. A. Lone, R. A. Leavitt, and D. A. Harrison, At. Data. Nucl. Data Tables **26**, 511 (1981).

[9] S. Raman, E. T. Jurney, J. W. Starner, and J. E. Lynn, Phys. Rev. **C 46**, 972 (1992).

[10] R. R. Roy, and B. P. Nigam, *Nuclear Physics*, (John Wiley Sons, inc. New York, 1967).

[11] P. M. Endt, Nucl. Phys. **A 521**, 1 (1990), *ibid.* **A 633**, 1 (1998).

[12] K. P. Lieb, H. G. Börner, M. S. Dewey, J. Jolie, S. J. Robinson, S. Ulbig, and Ch. Winter, Phys. Lett. **B 215**, 50 (1988).

[13] H. G. Börner and J. Jolie, J. Phys. G **19**, 217 (1993).



FIG. 1. Partial decay scheme for $^{29}$Si. The numbers in parentheses are the number of $\gamma$-rays per 100 neutron captures [8]. The spin and parity is referred to Ref. [9]



FIG. 2. Feynman diagram of neutron capture reaction $^{28}\mathrm{Si}(n, \gamma, \gamma)^{29}\mathrm{Si}$.



FIG. 3. The first part of the neutron capture reaction $^{28}\mathrm{Si}(n, \gamma, \gamma)^{29}\mathrm{Si}$ can be regarded as a neutron capture reaction $^{28}\mathrm{Si}(n, \gamma)^{29}\mathrm{Si}^*$ with emitting a single $\gamma$-ray.



FIG. 4. The second part of the neutron capture reaction $^{28}\mathrm{Si}(n, \gamma, \gamma)^{29}\mathrm{Si}$ can be regarded as a transition of an excited nucleus to the ground state with radiating a $\gamma$-ray .



FIG. 5. Schematic diagram of a coincidence measurement with considering the angular correlation of the two $\gamma$-rays.